

# HASTEN™

Adjuvant for improved performance of crop protection agents.

**Supplier SURfaPLUS Trading, Binnenhaven 1, 6709 PD Wageningen.  
Phone +31 317 451217 or e-mail [trading@surfaplus.com](mailto:trading@surfaplus.com)**

## **Prescription**

Allowed is only the application of this product as adjuvant/additive in combination with plant protection agents. Pay also attention to the label of the plant protection agent.

## **Direction for use**

Hasten™ contains esterified canola oil and an emulsifying system. Hasten™ improves the foliar uptake of several plant protection agents and improves the contact between insecticides and pests on the leaf surface. As a result, Hasten™ enhances the activity of several herbicides, insecticides, growth regulators and fungicides. Hasten reduces drift.

**Important** Each direction for use on the label of the plant protection agent has to be followed and has priority over the text on the label of Hasten™. Our recommendation is to apply the label rate of the plant protection agent. The use of Hasten™ with more than one plant protection agent in a mix or with plant protection agents not listed on this label is dissuaded without small scale tests in advance.

## **Applications**

### ***Herbicides***

Hasten™ can improve the performance of several herbicides including nicosulfuron, rimsulfuron, clodinafop-propargyl, quizalofop-P-ethyl, fluazifop-P-butyl, bentazone and isoproturon.

### ***Insecticides***

Hasten™ can improve the performance of several insecticides including abamectin, milbectin, spinosad, cyromazin and thiamethoxam.

### ***Fungicides***

Hasten™ can improve the performance of tebuconazole among others.

### ***Growth regulators***

Hasten™ can improve the performance of propiconazole and paclobutrazol.

Updates on the applications of Hasten™ can be found on [www.surfaplus-tr.com](http://www.surfaplus-tr.com) under Trading International.

### ***Crops***

Hasten™ can be applied in chrysanthemum, cucumbers, peppers and in arable farming. We recommend a small scale test in advance when Hasten is applied for the first time.

## **Preparation of spray solution**

If present, follow the instructions on the label of the crop protection agent. Our recommendation is to add Hasten™ as the last step in the preparation of the spray solution. Hasten™ mixes easily with the spray solution by gently stirring. Recommended concentration of Hasten™ is 250 ml per 100 liter (0.25% v/v). If preferred Hasten™ can be applied at 500 ml per 100 liter (0.5% v/v). Taking into account the various water volumes used in different crops, our advice is to include Hasten™ in such a way that the application rate of Hasten™ is between 0.2 and 2 L/ha.

**Storage**

Store well closed in original container at temperatures between 5 and 40 °C.

**This product is registered as adjuvant in the Netherlands under number 14376N**

**Hazard signal word: WARNING****Safety and risks**

Irritating to eyes.

Irritating to skin.

Wear protective gloves/protective clothing/eye protection/face protection.

In case of contact with eyes, rinse immediately with plenty of water and seek medical advice.

If on skin: wash with plenty of soap and water.

Keep out of reach of children.

Rinse the containers thoroughly with water before disposal.

*The information and the recommendations on this label are, to the best of our knowledge, correct on the date of packaging. Because we cannot control the storage, transport and the application of the product we do not guarantee the result. For this reason we do not accept responsibility for possible damage as a result of storage or transport conditions or as a result of the application. The sale of this product is subject to the conditions of sale of SURfaPLUS Trading.*